Cataract Eye Drop Schedule

Keep your eye closed after drop is instilled
and wait about 5 minutes between each drop.

DO NOT START DROPS UNTIL:_____

Before Surgery	3 Days Before	2 Days Before	1 Day Before
 Ofloxacin	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑
 Prednisolone	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑
 Diclofenac	❑❑❑	❑❑❑	❑❑❑

1st Week	Day of Surgery 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Ofloxacin	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑
Prednisolone	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑
Diclofenac	❑❑❑	❑❑❑	❑❑❑	❑❑❑	❑❑❑	❑❑❑	❑❑❑

2nd Week	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Ofloxacin	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑ 				
Prednisolone	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑	❑❑ ❑❑
Diclofenac	❑❑	❑❑	❑❑	❑❑	❑❑	❑❑	❑❑ 

3rd Week	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Prednisolone	❑❑❑	❑❑❑	❑❑❑	❑❑❑	❑❑❑	❑❑❑	❑❑❑

4th Week	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Prenisolone	❑❑	❑❑	❑❑	❑❑	❑❑	❑❑	❑❑ 